



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

*modramus sandwichensis bryanti* and *Geothlypis trichas sinuosa*; while from the southern marshes come the larger *Melospiza melodia cooperi*, *Ammodramus sandwichensis beldingi* and *Geothlypis trichas scirpicola*. It is also notable that the races of *Geothlypis trichas* occurring on the Pacific Coast are to a remarkable degree paralleled by those of the Atlantic Coast (See Palmer, Auk XVII, July 1900, pp. 216-242.)

In bringing together the material on which this paper is based, I am indebted for the loan of specimens to Messrs. F. S. Daggett, H. S. Swarth, W. O. Emerson and T. J. Hoover.



### Some Rare Birds in Los Angeles Co., Cal.

**P**HOEBE (*Sayornis phaebe*). On Feb. 14, 1901, I secured a male of this species near San Fernando. Not feeling certain as to its identity and having no specimens of my own with which to compare it, I sent it to Washington, where it was identified as *S. phaebe* by Dr. Chas. W. Richmond.

Slate-colored Sparrow. (*Passerella i. schistacea*). A female sparrow taken by myself at Millard's Canyon on Feb. 11, 1901 is identified, also by Dr. Richmond, as belonging to this subspecies. One other, also a female, taken by myself at Los Angeles and recorded by Grinnell in his 'List of Birds of the Pacific Slope of Los Angeles Co.' is I believe, the only one heretofore recorded from this part of the state. Another sparrow, a male, taken at Los Angeles Nov. 24, 1900 is described by the same authority as the last, as intermediate between *Passerella iliaca* and *P. i. schistacea*. This bird is rather sparsely spotted underneath, the throat and abdomen being immaculate, and the spots are of a brighter fox red than in the case of *P. i. schistacea* or *P. i. unalascensis*.

Flicker. (*Colaptes auratus*). I secured an adult female of this species at Los Angeles on Feb. 20, 1901. It was feeding in a pepper tree in company with several Red-shafted Flickers and was rather wild, in fact I had considerable difficulty in getting close enough to it to get a shot.

Besides these stragglers from the north and east, I have seen all through the past winter birds more or less com-

mon in the higher mountains, but rare or unknown at an altitude as low as the city of Los Angeles. On Nov. 14, 1900 I shot a female Williamson's Sapsucker (*Sphyrapicus thyroideus*); on Feb. 2, 1901 a female Townsend's Solitaire (*Myadestes townsendi*) and on Feb. 18 a male Red-naped Sapsucker (*Sphyrapicus varius nuchalis*). All these birds were feeding in pepper trees and within 50 yards of the same place. For the last two months Cassin's Purple Finch (*Carpodacus cassini*) has been quite abundant, feeding in pepper and willow trees. Nearly all that I have seen have had the plumage of the lower parts badly stained and gummed together, being often nearly black on the throat and breast.

Last week, April 15, I saw a flock of about two dozen Pine Siskins (*Spinus pinus*) feeding in some tall mustard, clinging to the top of the stems. I shot three and when I passed the spot several hours later the flock was still busily feeding. The three that I shot had their crops filled with plant lice, with which the wild mustard seed is covered. It struck me as rather an unusual diet for these birds. H. S. SWARTH.

Los Angeles, Cal., April 22, '01.



Harry R. Taylor made an interesting take on April 14 of White-tailed Kite  $\frac{1}{2}$ . We say "interesting," inasmuch as Mr. Taylor mentions that his pulse registered 140 beats per minute when he reached *terra firma!* Verily the spirit of olden days has not forsaken H. R. T!